


NNLO QCD CORRECTIONS TO $\bar{B} \rightarrow X_s \gamma$

M. MISIAK

*Institute of Theoretical Physics, Warsaw University,
Hoża 69, 00-681 Warsaw, Poland*



Current status of the NNLO QCD corrections to $\bar{B} \rightarrow X_s \gamma$ is reviewed. The calculations include three-loop matching conditions, four-loop anomalous dimensions as well as two- and three-loop on-shell amplitudes. Certain parts of the three-loop matrix elements are found by interpolation in the charm quark mass between the large- β_0 approximation in the $m_c = 0$ case and the complete result in the $m_c \gg m_b/2$ case.

1 Introduction

The weak radiative $\bar{B}$ -meson decay is known to be a sensitive probe of new physics. Thus, it is essential to calculate the Standard Model value of its branching ratio as precisely as possible. In order to do so, one writes

$$\begin{aligned} \mathcal{B}(\bar{B} \rightarrow X_s \gamma)_{E_\gamma > 1.6 \text{ GeV}} &= \mathcal{B}(\bar{B} \rightarrow X_c e \bar{\nu})^{\text{exp}} \left[\frac{\Gamma(b \rightarrow s \gamma)}{\Gamma(b \rightarrow c e \bar{\nu})} \right]_{\text{LO EW}} f \left(\frac{\alpha_s(M_W)}{\alpha_s(m_b)} \right) \times \\ &\times \left\{ 1 + \underbrace{\mathcal{O}(\alpha_s)}_{\sim 25\%} + \underbrace{\mathcal{O}(\alpha_s^2)}_{\sim 7\%} + \underbrace{\mathcal{O}(\alpha_{\text{em}})}_{\sim 4\%} + \underbrace{\mathcal{O}\left(\frac{\Lambda^2}{m_b^2}\right)}_{\sim 1\%} + \underbrace{\mathcal{O}\left(\frac{\Lambda^2}{m_c^2}\right)}_{\sim 3\%} + \underbrace{\mathcal{O}\left(\frac{\alpha_s \Lambda}{m_b}\right)}_{< \sim 5\%} \right\}, \end{aligned} \quad (1)$$

where $\mathcal{B}(\bar{B} \rightarrow X_c e \bar{\nu})^{\text{exp}}$ is the measured semileptonic branching ratio. The b -quark radiative and semileptonic decay widths $\Gamma(b \rightarrow s \gamma)$ and $\Gamma(b \rightarrow c e \bar{\nu})$ are calculated perturbatively at the leading order in electroweak interactions and neglecting QCD effects. Normalization to the semileptonic rate is introduced to eliminate uncertainties from the CKM angles and overall factors of m_b^5 . The Leading Order (LO) QCD correction factor is denoted by $f(\alpha_s(M_W)/\alpha_s(m_b))$.

Higher-order perturbative and non-perturbative corrections are listed in the second line of Eq. 1. Their approximate sizes are indicated. The non-perturbative effects arise only as corrections, thanks to the heaviness of the b -quark ($m_b \gg \Lambda \equiv \Lambda_{\text{QCD}}$) and to the inclusive character

of the considered decays.¹ The $\mathcal{O}(\alpha_s)$, $\mathcal{O}(\alpha_{\text{em}})$, $\mathcal{O}(\Lambda^2/m_b^2)$ and $\mathcal{O}(\Lambda^2/m_c^2)$ contributions are known since many years.² On the other hand, the indicated sizes of the $\mathcal{O}(\alpha_s^2)$ and $\mathcal{O}(\alpha_s\Lambda/m_b)$ corrections are only estimates that were made before the actual calculation of the Next-to-Next-to-Leading (NNLO) QCD corrections ($\mathcal{O}(\alpha_s^2)$).³

The current experimental world average for the branching ratio reads⁴

$$\mathcal{B}(\bar{B} \rightarrow X_s \gamma)_{E_\gamma > 1.6 \text{ GeV}}^{\text{exp}} = \left(3.55 \pm 0.24 \begin{smallmatrix} +0.09 \\ -0.10 \end{smallmatrix} \pm 0.03\right) \times 10^{-4}, \quad (2)$$

where the combined error is approximately equal to the expected size of the $\mathcal{O}(\alpha_s^2)$ effects. Thus, the currently reported NNLO calculation is well-motivated. It is hoped that the $\mathcal{O}(\alpha_s\Lambda/m_b)$ uncertainty can be reduced in the future by performing a dedicated analysis.




Figure 1: Sample 1-loop diagram.




Figure 2: Sample 2-loop diagram.

Fig. 1 presents a sample leading-order electroweak diagram for $b \rightarrow s\gamma$ in the SM. Two-loop diagrams like the one shown in Fig. 2 contain large logarithms $\ln M_W^2/m_b^2$ which enhance the branching ratio by more than a factor of two. Such logarithms are resummed from all orders of the perturbation series into the LO QCD correction factor $f(\alpha_s(M_W)/\alpha_s(m_b))$. Resummation of large logarithms is necessary at higher orders, too. Since two loops are relevant at the LO, four loops are necessary at the NNLO, which makes the $\mathcal{O}(\alpha_s^2)$ calculation extremely involved.

2 The effective Lagrangian

Resummation of $(\alpha_s \ln M_W^2/m_b^2)^n$ at each order in α_s is most conveniently performed in the framework of an effective theory that arises from the SM after decoupling of the heavy electroweak bosons and the top quark. The Lagrangian of such a theory reads

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{QCD} \times \text{QED}}(u, d, s, c, b) + \frac{4G_F}{\sqrt{2}} V_{ts}^* V_{tb} \sum_{i=1}^8 C_i(\mu) Q_i. \quad (3)$$

The operators Q_i and numerical values of their Wilson coefficients at $\mu \simeq m_b$ are as follows:

$$Q_i = \begin{cases} (\bar{s}\Gamma_i c)(\bar{c}\Gamma'_i b), & i = 1, 2, & |C_i(m_b)| \sim 1, \\ (\bar{s}\Gamma_i b)\Sigma_q(\bar{q}\Gamma'_i q), & i = 3, 4, 5, 6, & |C_i(m_b)| < 0.07, \\ \frac{em_b}{16\pi^2} \bar{s}L\sigma^{\mu\nu} b_R F_{\mu\nu}, & i = 7, & C_7(m_b) \sim -0.3, \\ \frac{gm_b}{16\pi^2} \bar{s}L\sigma^{\mu\nu} T^a b_R G_{\mu\nu}^a, & i = 8, & C_8(m_b) \sim -0.15. \end{cases} \quad (4)$$

Here, Γ and Γ' stand for various products of the Dirac and color matrices.⁶ The perturbative calculations are performed in three steps: **(i) Matching:** Evaluating $C_i(\mu_0)$ at $\mu_0 \sim M_W$ by requiring equality of the SM and effective theory Green's functions at the leading order in (external momenta)/ M_W . **(ii) Mixing:** Deriving the effective theory Renormalization Group Equations (RGE) and evolving $C_i(\mu)$ from μ_0 down to $\mu_b \sim m_b$. **(iii) Matrix elements:** Evaluating the on-shell $b \rightarrow X_s^{\text{parton}} \gamma$ amplitudes at $\mu_b \sim m_b$.




Figure 3: Corrections δ_1 , $\delta_2^{\beta_0}$, δ_2^{rem} and δ_3 ($\times 100$) as functions of $r = m_c(m_c)/m_b^{1S}$ (see the text).

3 Current status of the NNLO calculation

The 3-loop matching for Q_7 and Q_8 was evaluated more than two years ago.⁵ The 3-loop mixings in the Q_1 - Q_6 and Q_7 - Q_8 sectors were found more recently.^{6,7} The yet unpublished results⁸ on the 4-loop mixing of Q_1 - Q_6 into Q_7 will be used below. The effect of the unknown 4-loop mixing of Q_1 - Q_6 into Q_8 is expected to be small. Nevertheless, its calculation is underway.⁸

As far as the matrix elements are concerned, contributions to the decay rate that are proportional to $|C_7(\mu_b)|^2$ are completely known at the NNLO.^{9,10} These two-loop results have recently been confirmed by independent groups.^{11,12}

Two- and three-loop matrix elements in the so-called large- β_0 approximation were found¹³ as an expansion in m_c/m_b that is convergent for $m_c < m_b/2$, i.e. in the physical domain. The remaining (“beyond-large- β_0 ”) contributions to the matrix elements were calculated¹⁴ in the limit $m_c \gg m_b/2$ and then interpolated to smaller values of m_c (see the next section).

In order to relate our result with $E_{\text{cut}} = 1.6 \text{ GeV}$ to the measurements with cuts at 1.8 GeV (Belle¹⁵) and 1.9 GeV (BaBar¹⁶), one needs to evaluate ratios of the decay rates with different cuts. Such a calculation at the NNLO has recently been completed.¹⁷ However, the final numerical results are not yet available, and the average in Eq. 2 does not include them.

4 Interpolation in m_c

Let us parametrize the NNLO correction to the branching ratio in terms of three quantities δ_i

$$\mathcal{B}(\bar{B} \rightarrow X_s \gamma)_{E_\gamma > 1.6 \text{ GeV}} \equiv \mathcal{B}_{\text{NNLO}}(r) = \mathcal{B}_{\text{NLO}}(r) + \mathcal{B}_{\text{NLO}}(0.262) (\delta_1 + \delta_2(r) + \delta_3(r)), \quad (5)$$

where $r = m_c(m_c)/m_b^{1S}$ and $\mathcal{B}_{\text{NLO}}(0.262) \simeq 3.38 \times 10^{-4}$. The quantities δ_i contain terms depending on different contributions to the Wilson coefficient perturbative expansion

$$C_i(\mu_b) = C_i^{(0)}(\mu_b) + \frac{\alpha_s(\mu_b)}{4\pi} C_i^{(1)}(\mu_b) + \left(\frac{\alpha_s(\mu_b)}{4\pi} \right)^2 C_i^{(2)}(\mu_b) + \dots \quad (6)$$

In particular, δ_1 contains terms proportional to $C_i^{(0)} C_j^{(2)}$ and $C_i^{(1)} C_j^{(1)}$, δ_2 – terms proportional to $C_i^{(0)} C_j^{(0)}$, and δ_3 – terms proportional to $C_i^{(0)} C_j^{(1)}$. The so-called large- β_0 part of δ_2 is found from the fermionic contributions to this quantity: $\delta_2 = A n_f + B = \delta_2^{\beta_0} + \delta_2^{\text{rem}}$, where $\delta_2^{\beta_0} = -\frac{3}{2}(11 - 2/3 n_f)A$ and $\delta_2^{\text{rem}} = \frac{33}{2}A + B$. Here, $n_f = 5$ is the number of active flavors in the effective theory. While $\delta_2^{\beta_0}$ is known^{13,14} for all r , δ_2^{rem} has been calculated¹⁴ only for $r \gg \frac{1}{2}$ and needs to be interpolated to lower values of r .

Preliminary results of this interpolation are summarized in Fig. 3 (for $\mu_b = m_b$). The two vertical lines mark the experimental bounds on r . Dashed curves show the calculated asymptotic

behavior of each δ_i for $r \gg \frac{1}{2}$ at the leading order in $1/(4r^2)$, i.e. including only the constant and logarithmic terms. Solid lines show either the known exact dependence on r (for δ_1 and δ_3), the known small- r expansion (for $\delta_2^{\beta_0}$) or the interpolation (for δ_2^{rem}). The interpolation can be performed approximating δ_2^{rem} by a linear combination of four functions:

$$\delta_2^{\text{rem}}(r) = a \mathcal{B}_{\text{NLO}}(r) + b r \frac{d}{dr} \mathcal{B}_{\text{NLO}}(r) + c \delta_2^{\beta_0}(r) + d \quad (7)$$

The coefficients a , b , c and d are determined in a unique manner from the asymptotic behavior at large r and from the requirement that either $\delta_2^{\text{rem}}(0) = 0$ (lower curves) or $\delta_1 + \delta_2^{\text{rem}}(0) + \delta_3(0) = 0$ (upper curves). The assumed functional dependence of δ_2^{rem} on r is motivated by the r -dependence of renormalization-induced effects. In many explicitly calculated examples, such effects have been found to dominate over other terms of the same order.

5 Conclusion

The two ways of interpolation¹⁴ lead to two values of the NNLO branching ratio $\mathcal{B}_{\text{NNLO}} = 3.06 \times 10^{-4}$ or $\mathcal{B}_{\text{NNLO}} = 3.24 \times 10^{-4}$, for $E_\gamma > 1.6 \text{ GeV}$. These values are around 6% apart from each other and around 7% below the NLO result $\mathcal{B}_{\text{NLO}} = 3.38 \times 10^{-4}$. Their average 3.15×10^{-4} is around $1.5\sigma_{\text{exp}}$ below the experimental result in Eq. 2. Consequently, extensions of the SM predicting a suppression of the $b \rightarrow s\gamma$ amplitude are going to be more constrained than previously.

6 Acknowledgments

I am greatly indebted to the contributors to Refs. 5,6,7,8,9,10,11,12,13,14 for their participation in the NNLO enterprise. I am grateful to the organizers of the 40th Rencontres de Moriond for their invitation and care. I acknowledge support from the Polish Committee for Scientific Research under the grant 2 P03B 078 26 and from the European Community's Human Potential Programme under the contract HPRN-CT-2002-00311, EURIDICE.

References

1. J. Chay, H. Georgi and B. Grinstein, *Phys. Lett. B* **247**, 399 (1990).
2. A.J. Buras and M. Misiak, *Acta Phys. Pol. B* **33**, 2597 (2002).
3. M. Misiak *et al.*, hep-ph/0609232.
4. T. Barberio *et al.* (HFAG), hep-ex/0603003.
5. M. Misiak and M. Steinhauser, *Nucl. Phys. B* **683**, 277 (2004).
6. M. Gorbahn and U. Haisch, *Nucl. Phys. B* **713**, 291 (2005).
7. M. Gorbahn, U. Haisch and M. Misiak, *Phys. Rev. Lett.* **95**, 102004 (2005).
8. M. Czakon, U. Haisch and M. Misiak, unpublished.
9. I. Blokland *et al.*, *Phys. Rev. D* **72**, 033014 (2005).
10. K. Melnikov and A. Mitov, *Phys. Lett. B* **620**, 69 (2005).
11. H. M. Asatrian *et al.*, *Nucl. Phys. B* **749**, 325 (2006).
12. H. M. Asatrian, T. Ewerth, A. Ferroglia, P. Gambino and C. Greub, hep-ph/0607316.
13. K. Bieri, C. Greub and M. Steinhauser, *Phys. Rev. D* **67**, 114019 (2003).
14. M. Misiak and M. Steinhauser, hep-ph/0609241.
15. P. Koppenburg *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **93**, 061803 (2004).
16. B. Aubert *et al.* (BaBar Collaboration), hep-ex/0507001.
17. T. Becher and M. Neubert, *Phys. Lett. B* **633**, 739 (2006), *Phys. Lett. B* **637**, 251 (2006).